TRANSGÉNICOS

Olvido Périz Ponce

Verónica Pedroza Peral

TÉCNICAS PARA LA PRODUCCIÓN DE ANIMALES TRANSGÉNICOS

Transferencia de genes mediante vectores retrovirales.

TÉCNICAS PARA LA PRODUCCIÓN DE ANIMALES TRANSGÉNICOS

Producción de quimeras transgénicas a partir de células y embriones genéticamente modificados.

TÉCNICAS PARA LA PRODUCCIÓN DE ANIMALES TRANSGÉNICOS

Microinyección de ADN en el pronúcleo del cigoto.

TÉCNICAS PARA LA PRODUCCIÓN DE PLANTAS TRANSGÉNICAS

PLANTAS TRANSGÉNICAS

GLOBAL AREA OF BIOTECH CROPS Million Hectares (1996-2012)

A record 17.3 million farmers, in 28 countries, planted 170.3 million hectares (420 million acres) in 2012, a sustained increase of 6% or 10.3 million hectares (25 million acres) over 2011.

Source: Clive James, 2012.

www.isaaa.org

Situación mundial de los CULTIVOS BIOTECNOLÓGICOS en el 2011

Fuente: Clive James 2012. ISAAA Brief 43-2011.

Cultivos transgénicos en el mundo

Cultivos biotecnológicos en el mundo
(millones de hectáreas)

Global Adoption Rates (%) for Principal Biotech Crops (Million Hectares, Million Acres), 2012

M Acres

Source: Clive James, 2012

ESTIMACIÓN DE LA SUPERFICIE TOTAL DE VARIEDADES OMG CULTIVADAS EN ESPAÑA

COMUNIDAD AUTÓNOMA	SUPERFICIE (ha)
Andalucía	10.361
Aragón	41.669
Castilla La Mancha	7.883
Castilla León	8
Catalunya	33.530
Madrid	420
Navarra	5.800
Valencia	522
Extremadura	15.951
Baleares	154
Murcia	4
TOTAL	116.302

España tiene 17 millones de hectáreas de terreno cultivable, de los cuales 369.000 están dedicadas al cultivo de las distintas variedades de maíz (un 2,1% del total).

Dentro de los cultivos dedicados al maíz, un 26,4% usa la modalidad OMG MON 810. Este dato está calculado en función de las semillas de MON 810 vendidas en cada región, a razón de 85.000 semillas por hectárea.

La estimación de superficie está referida a una dosis media de 85.000 semillas/ha

Aplicaciones

- Tolerancia a herbicidas
- Resistencia a insectos
- Resistencia a patógenos (sobre todo virus)
- Resistencia a estrés abiótico
 - Resistencia a sequía (maíz)
 - Trabajando en tolerancia a salinidad, calor, etc.
- Mejora de la calidad nutritiva
- Biorremediación
 - Remediación de suelos contaminados
- Biorreactores
 - Medicamentos
 - Vacunas

Resistencia a herbicidas

Soja tolerante a glifosfato (derecha) y sin tratar con el herbicida (izquierda)

Resistencia a insectos

Genes Cry	Tóxicos para
CryIA(a), CryIA(b), CryIA(c)	Lepidóptera
Cry1B, Cry1C, Cry1D	Lepidóptera
CryII	Lepidóptera, Diptera
CryIII	Coleóptera
CryIV	Diptera
CryV	Lepidóptera, Coleóptera

Maíz atacado por barrenador del tallo

Trigo para celíacos

Equipo de investigación del CSIC a cargo de la investigación del trigo apto para celíacos

Arroz dorado

Mapa de la incidencia mundial de la deficiencia en vitamina A

Arroz dorado

Arroz convencional, Arroz dorado I y Arroz dorado II

Esquema de enzimas clonadas

Arroz dorado

Biorreactores: vacunas comestibles

- Vacunas de tomate contra SIDA y hepatitis B (probadas en animales).
- Patata contra la hepatitis B y enfermedad de Newcastle (ensayos exitosos en humanos).
- Plátano contra la hepatitis B (en desarrollo).
- Alfalfa contra la fiebre aftosa para el ganado bovino (en estudio).

Vacunas comestibles

Mazorcas de maíz y mazorcas de maíz expresando un fármaco

Otras aplicaciones

► Frutos más resistentes

ARN antisentido inhibe la producción de la enzima poligalacturonasa

► Tolerancia a estrés abiótico

USO DE LOS ANIMALES TRANSGÉNICOS

1. Ayudar en la identificación, aislamiento y caracterización de genes y secuencias importantes para la expresión génica.

USO DE LOS ANIMALES TRANSGÉNICOS

2. Generar modelos de enfermedades que afectan al hombre, para el desarrollo de nuevas drogas y tratamientos.

Ratones knock-out:

- Bajo costo.
- Rápido periodo de gestación.
- Facilidad de manejo.

Sin embargo... POCA RELACIÓN
FILOGENÉTICA CON LOS
HUMANOS.

Ratones knock-out en el estudio de enfermedades neurodegenerativas.

Instituto Nacional de Investigación
y Tecnología Agraria y Alimentaria

USO DE LOS ANIMALES TRANSGÉNICOS

3. Animales transgénicos como biorreactores. Producir moléculas de interés industrial.

USO DE LOS ANIMALES TRANSGÉNICOS

4. Fuente de tejidos y órganos para trasplantes en humanos.

Xenotrasplantes → Aún en estudio, debido a:

- Rechazo hiper-agudo.
- Rechazo vascular retardado.
- Rechazo crónico de injertos.
- Rechazo celular.

USO DE LOS ANIMALES TRANSGÉNICOS

5. Mejorar el ganado y otros animales de importancia económica.

“Súper salmón”, el posible primer animal transgénico que el ser humano pueda degustar.

Vacas frisonas transgénicas contra la tuberculosis bovina

CONTROVERSIA

- Reducción de la biodiversidad
- Proceso "antinatural"
- Patentes
- Medios de comunicación
- Seguridad
- Ética del uso de animales transgénicos
- Etiquetado

La transgénesis es “antinatural”

El problema de las patentes

- Existen transgénicos libres de patentes como el arroz dorado.
- Países como Brasil, Argentina, India y China liberan las patentes para sus agricultores.
- Las multinacionales son las únicas capaces de costear los controles exigidos.

EVENTOS BIOTECNOLÓGICOS AUTORIZADOS EN LA UNIÓN EUROPEA

La UE tiene **50** eventos biotecnológicos aprobados:
48 para importación y sólo dos para cultivo.

48

2

Resumen proceso autorización OMGs en la UE

Autorizados para importación (45)

Autorizados para cultivo (2)

- 1** MAÍZ Bt

Maíz resistente a la plaga del taladro. España es el país europeo con más superficie cultivada: 131.538 has. en 2013.
- 1** PATATA AMFLORA

Patata que produce un almidón de amilopectina pura, idóneo para aplicaciones técnicas. Pese a estar aprobado su cultivo en la UE no se comercializa a día de hoy.

En proceso de autorización (68)

ALIMENTACIÓN, PROCESADO, PIENSOS E IMPORTACIÓN
maíz (16), algodón (12), soja (11), colza (6), patata (1), arroz (1)

47

CULTIVO
maíz (17), patata (2), soja (1), remolacha azucarera (1)

21

Retrasos en autorizaciones de OMGs

Más de **44** años de retrasos acumulados en autorización de eventos MG a 1 de junio de 2014.

El caso del maíz 1507

Lleva 12 años esperando ser aprobado y ya ha recibido siete opiniones científicas favorables de seguridad por parte de la EFSA.

Medios de comunicación

El príncipe Carlos y su esposa. (Foto: REUTERS)

POR EL ENDEUDAMIENTO DE LOS AGRICULTORES

Carlos culpa a los transgénicos de los suicidios en la India

El príncipe de Gales se refirió al "número horrible de suicidios de pequeños agricultores en la India debidos parcialmente al fracaso de las variedades de cosechas transgénicas".

Aunque los suicidios se produjeron antes de la introducción de las cosechas transgénicas en ese país, algunos ecologistas creen que los agricultores se vieron forzados a comprar las nuevas semillas, que no sólo son más caras, sino que además no produjeron cosechas significativas.

A STEADY RATE OF TRAGEDY

Contrary to popular myth, the introduction in 2002 of genetically modified Bt cotton is not associated with a rise in suicide rates among Indian farmers.

Seguridad

- Principio de equivalencia sustancial (PES)
- Efectos imprevistos: pleitropía
- Oposición
- Resistencia a antibióticos
- Alergias
- Transferencia horizontal de genes

Gilles-Eric Seralini

1. Introduction

There is an ongoing concern about the necessary weight of scientific evidence in relation to the acceptance of genetically modified organisms (GMOs), including regular toxicologic and epidemiologic studies, the regulatory authority of the industry, animal testing studies to be performed, and the potential for unintended genetic effects, among others, of Roundup and Roundup-tolerant crops. These investigations mostly concern GM and maize that are combined either herbicide-tolerant (to Roundup) or in 80% of cases, or engineered to produce a modified Bt toxin (Bt-toxin), or both. As a result, these GM crops contain two pesticide herbicides for which one maximal residual level (MRL) has been established in some countries. The pesticides combine to generate that there is no major change in genetically modified organisms (GMO) herbicides, according to studies (Shenoy and Girdhar, 2011; Hammond et al., 2004, 2006a,b). Significant disturbances have been found and may be interpreted differently (Seralini et al., 2005; Seralini de Vendômois et al., 2010). Detailed analyses have revealed alterations in kidney and liver functions that may be the signs of early chronic liver intoxication, possibly explained at least in part by pesticide residues in the GM food (Seralini et al., 2007; Seralini de Vendômois et al., 2009; Seralini, 2010). Seralini de Vendômois et al. (2010) have also demonstrated that 8 concentrations in the range of 10³ times below the MRL induced endocrine disturbances in human cells (Seralini et al., 2005) and toxic effects (Seralini, Seralini de Vendômois and Seralini, 2009), including in vivo (Seralini et al., 2011). After several months of administration of an 8 herbicide mix, the liver and pancreas of rats were affected, as highlighted by disturbances in both nuclear structure (Seralini et al., 2006a, 2006b). Furthermore, this toxic effect was reproduced by the application of 8 herbicide already in September in culture (Seralini et al., 2008).

Abbreviations: USA generally available; 2 Roundup MRL, maximal residual level; USA, generally available; 80% Bt-toxin, transgenic Bt-toxin; 8 herbicide-tolerant (to Roundup) or in 80% of cases, or engineered to produce a modified Bt toxin (Bt-toxin), or both; GM crops contain two pesticide herbicides for which one maximal residual level (MRL) has been established in some countries; The pesticides combine to generate that there is no major change in genetically modified organisms (GMO) herbicides, according to studies (Shenoy and Girdhar, 2011; Hammond et al., 2004, 2006a,b); Significant disturbances have been found and may be interpreted differently (Seralini et al., 2005; Seralini de Vendômois et al., 2010); Seralini de Vendômois et al. (2010) have also demonstrated that 8 concentrations in the range of 10³ times below the MRL induced endocrine disturbances in human cells (Seralini et al., 2005) and toxic effects (Seralini, Seralini de Vendômois and Seralini, 2009), including in vivo (Seralini et al., 2011). After several months of administration of an 8 herbicide mix, the liver and pancreas of rats were affected, as highlighted by disturbances in both nuclear structure (Seralini et al., 2006a, 2006b). Furthermore, this toxic effect was reproduced by the application of 8 herbicide already in September in culture (Seralini et al., 2008).

© 2011 Elsevier Ltd. All rights reserved.
doi:10.1016/j.fct.2011.08.004

MANAGING EDITORS:

BRYAN DELANEY – ANKENY, IA
JOSÉ L. DOMINGO – CATALONIA, SPAIN
SIEGFRIED KNASMUELLER – VIENNA, AUSTRIA

REVIEW EDITOR:

SUSAN BARLOW – BRIGHTON, U.K.

ASSOCIATE EDITORS:

SILVIA BERLANGA DE M BARROS – SÃO PAULO, BRAZIL
DAVID J. BRUSICK – BUMPASS, VA
ADRIAN COVACI – WILRIJK-ANTWERPEN, BELGIUM
WILLIAM C. HALL – MOUNT AIRY, MD
SALMAAN INAYAT-HUSSAIN – KUALA LUMPUR, MALAYSIA
CLAIRE L. KRUGER – BETHESDA, MD
JOHN CHRISTIAN LARSEN – SØBORG, DENMARK
BYUNG-MU LEE – SUWON, GYEONGGI-DO, SOUTH KOREA
JUAN M. LLOBET – BARCELONA, SPAIN
MICHAEL W. PARIZA – MADISON, WI
IVONNE M.C.M. RIETJENS – WAGENINGEN, NETHERLANDS
SHUJI TSUDA – MORIOKA, IWATE, JAPAN
JEAN-LUC VOLATIER – MAISONS ALFORT, FRANCE
GARY WILLIAMS – VALHALLA, NY
PAUL WRIGHT – BUNDOORA, AUSTRALIA
PING-KUN ZHOU – BEIJING, CHINA

November 19, 2013

Professor G. E. Séralini
University of Caen
Institute of Biology, EA 2608
CRIIGEN and Risk Pole CNRS
Esplanade de la Paix
14032 Caen Cedex FRANCE

Dear Professor Séralini,

As you know, several months ago, we began a thorough examination of the data you provided to the journal. The panel looking at the data and its analysis sought to address the questions that had been raised in the Letters to the Editor received in response to your article.

The panel had many concerns about the quality of the data, and ultimately recommended that the article should be withdrawn. I have been trying to get in touch with you to discuss the specific reasons behind this recommendation. If you do not agree to withdraw the article, it will be retracted, and the following statement will be published in its place:

The journal Food and Chemical Toxicology retracts the article “Long term toxicity of a Roundup herbicide and a Roundup-tolerant genetically modified maize,”¹ which was published in this journal in November 2012. This retraction comes after a thorough and time-consuming analysis of the published article and the data it reports, along with an investigation into the peer-review behind the article. The Editor in-Chief deferred making any public statements regarding this article until this investigation was complete, and the authors were notified of the findings.

Very shortly after the publication of this article, the journal received Letters to the Editor expressing concerns about the validity of the findings it described, the proper use of animals,

and even allegations of fraud. Many of these letters called upon the editors of the journal to retract the paper. According to the journal's standard practice, these letters, as well as the letters in support of the findings, were published along with a response from the authors.² Due to the nature of the concerns raised about this paper, the Editor-in-Chief examined all aspects of the peer review process and requested permission from the corresponding author to review the raw data. The request to view raw data is not often made; however, it is in accordance with the journal's policy that authors of submitted manuscripts must be willing to provide the original data if so requested.³ The corresponding author agreed and supplied all material that was requested by the Editor-in-Chief. The Editor-in-Chief wishes to acknowledge the co-operation of the corresponding author in this matter, and commends him for his commitment to the scientific process.

Unequivocally, the Editor-in-Chief found no evidence of fraud or intentional misrepresentation of the data. However, there is legitimate cause for concern regarding both the number of animals in each study group and the particular strain selected. The low number of animals had been identified as a cause for concern during the initial review process, but the peer review decision ultimately weighed that the work still had merit despite this limitation. A more in-depth look at the raw data revealed that no definitive conclusions can be reached with this small sample size regarding the role of either NK603 or glyphosate in regards to overall mortality or tumor incidence. Given the known high incidence of tumors in the Sprague-Dawley rat, normal variability cannot be excluded as the cause of the higher mortality and incidence observed in the treated groups.

Ultimately, the results presented (while not incorrect) are inconclusive, and therefore do not reach the threshold of publication for Food and Chemical Toxicology. The peer review process is not perfect, but it does work. The journal is committed to a fair, thorough, and timely peer-review process; sometimes expediency might be sacrificed in order to be as thorough as possible. The time-consuming nature is, at times, required in fairness to both the authors and readers. Likewise, the Letters to the Editor, both pro and con, serve as a post-publication peer-review. The back and forth between the readers and the author has a useful and valuable place in our scientific dialog.

The Editor-in-Chief again commends the corresponding author for his willingness and openness in participating in this dialog. The retraction is only on the inconclusiveness of this one paper. The journal's editorial policy will continue to review all manuscripts no matter how controversial they may be. The editorial board will continue to use this case as a reminder to be as diligent as possible in the peer review process.

Sincerely,

A. Wallace Hayes
Editor-In-Chief

BIOÉTICA EN LA PRODUCCIÓN Y USO DE ANIMALES TRANSGÉNICOS

Manipulación genética animal = Problemas éticos

BIOÉTICA EN LA PRODUCCIÓN Y USO DE ANIMALES TRANSGÉNICOS

En el 10% de los casos de transgénesis en animales se generan efectos inesperados e impredecibles.

¿El avance del conocimiento es razón suficiente para considerar como sacrificable la vida de los animales de laboratorio?

BIOÉTICA EN LA PRODUCCIÓN Y USO DE ANIMALES TRANSGÉNICOS

Personas = Sujetos morales

Animales = Objetos morales

¿EL SUFRIMIENTO DE OTRAS FORMAS DE VIDA ES NECESARIO PARA UNA MEJORA EN LA VIDA HUMANA?

BIOÉTICA EN LA PRODUCCIÓN Y USO DE ANIMALES TRANSGÉNICOS

Principio de las tres R's, de Russel y Burch (1959):

Reducción.

Refinamiento.

Reemplazo.

ETIQUETADO DE LOS ALIMENTOS TRANSGÉNICOS

Los consumidores tienen derecho a elegir libremente y al buen conocimiento de lo que consumen. Para ello se utiliza el etiquetado de los alimentos.

ETIQUETADO DE LOS ALIMENTOS TRANSGÉNICOS

“PROPOSICIÓN 37”